

# MISSOURI MONTHLY VITAL STATISTICS

## *Provisional Statistics*

FROM THE

MISSOURI DEPARTMENT OF HEALTH

CENTER FOR HEALTH INFORMATION MANAGEMENT & EPIDEMIOLOGY

Jefferson City, Missouri 65102-0570

(573) 751-6272

October 1998 VOL. 32 NO. 8

### *Focus. . . Alzheimer's Disease Mortality*

The number of deaths attributed to Alzheimer's disease in Missouri and nationally has increased markedly over recent years. (See Figure 1.) The number of Missouri resident Alzheimer's disease deaths in 1996 was 14.6 times the 1980 number; nationally, the increase was more than sixteenfold. If Alzheimer's disease had been counted separately as a leading cause of death in Missouri, it would have ranked eleventh in 1996 and tenth in 1997. Nationally, Alzheimer's disease moved from fourteenth in 1995 to thirteenth in 1996.

A study of the records from death certificates can tell us about patterns in this increase, and it provides explanations for some of the increase.


Alzheimer's disease (AD) is best known as a disease that progressively destroys memory. It acts both to destroy brain cells and to interrupt the communication among cells. Patients are not only forgetful and confused but may also have prob-

lems with judgment, delusions, unusual behaviors, and unstable moods. Eventually, patients become unable to care for themselves and are likely to become bedridden. They are increasingly vulnerable to infectious diseases; pneumonia is listed as having contributed to about seventeen percent of Missouri's AD deaths.

Alzheimer's disease is named for Alois Alzheimer, a German doctor who in 1906 discovered abnormal clumps and tangled bundles of fibers in the brain of a woman who had died of a dementia. Those plaques and tangles are now considered the definitive diagnostic signs of AD.<sup>1</sup>

AD is primarily a disease of the elderly. Of Missouri resident AD deaths in 1997, half (50.4 percent) were of persons over 85, and 99.0 percent were of persons over age 65. In that year, AD accounted for less than one percent (0.9) of all Missouri resident deaths, but 1.2 percent of deaths over age 65,

**Figure 1**  
**Missouri Resident Deaths Attributed to Alzheimer's Disease 1980-1997**



and 1.6 percent of deaths over age 85.

Because AD is primarily a disease of the elderly, the fact that our population is aging is one explanation to consider as the cause of the increase in AD deaths. As death rates from other chronic diseases such as heart disease and cancer decline, more and more people live long enough to develop AD. Also, some persons who might have died of another disease while in early stages of AD now survive the other disease and die of AD.

In 1980, persons 85 years and older represented 1.2 percent of Missouri's population; in 1990, they were 1.6 percent, and estimates for 1997 are that they were 1.8 percent. The percent 75 and older went up from 5.4 percent in 1980 to an estimated 6.5 percent in 1997. Those increases are too small to explain much of the fourteenfold increase in AD deaths in Missouri between 1980 and 1996.

Figure 2 shows that the age group 85 and older has not only the largest rate of AD deaths, but also the largest increase in the AD death rate per 100,000 population. In the oldest age group, the AD death rate more than doubled between 1987 and 1997, from 112.2 to 262.7 (134.1 percent). The rate for persons 65 to 74 does not show a consistent trend over the past decade, while for ages 75 to 84 it increased 51.3 percent, from 49.1 to 74.3 per 100,000 population. For whatever reason, the increase in AD deaths is concentrated in the oldest of the elderly.

Figure 2 provides the answer to another question, as well: the role of gender. Since the beginning of 1980, 3,054 of the 4,791 Missouri residents whose deaths were attributed to AD have been female. That is 63.7 percent; a similar percentage of the deaths attributed to senility and other dementias were female. Does this reflect a higher death rate due to AD and similar conditions for females, or just that they are more likely

to survive into the years of high risk for those conditions? Figure 2 shows that age is a far more important factor than gender, and females do not have consistently higher age-specific AD death rates. Indeed, national statistics showed a higher rate for males during the 1980's.<sup>2</sup>


Another explanation to be considered is that doctors are becoming more likely to supply "Alzheimer's disease" as the cause of death in cases of dementia. As AD has become better known, doctors may also be more aware of it. Diagnostic tools are improving. Also, as certifying physicians who use vaguer terms such as "senility" are asked for more specific information, they may become more likely to supply "Alzheimer's disease."


However, Figure 3 shows that the number of deaths attributed to similar conditions (senility, presenile dementias and other cerebral degenerations) also increased sharply over the same time period. Clearly, the number of attributions of death to AD was not merely subtracted from those that otherwise would have been attributed to other dementias. However, there may be an increase in attributions to Alzheimer's disease for deaths that in earlier years would have been attributed to other kinds of diseases.

Although aging of the population and increased reporting may be contributing to the increase in AD deaths, there still may be other unrecognized reasons for the increase in deaths due to dementia in the elderly population.

Any increase in dementias among the elderly has implications for the future. The Bureau of the Census estimates that the number of Americans over age 85 will grow from about 4.4 million in the year 2001 to seven million in 2025 and 8.5 million in 2030.<sup>3</sup> Even without an increase in age-specific rates, an increasing number of elderly citizens leads to predic

**Figure 2**  
**Alzheimer's Disease Death Rates by Age and Gender**  
**Missouri Residents 1987 - 1997**





## Provisional Vital Statistics for August 1998

**Live births** increased in August as 5,908 babies were born compared with 5,750 one year earlier. Because of a longer reporting period in 1998, the rate actually decreased slightly during this time period from 12.5 to 12.4 per 1,000 population.

Cumulative births decreased for the 8 months ending with August, but increased for the 12 months ending with August.

**Deaths** increased in August, but decreased for the cumulative 8- and 12-month periods ending with August. Deaths decreased by 3.5 percent for the first eight months on 1998.

The **Natural increase** for Missouri in August was 1,378

(5,908 births minus 4,530 deaths) compared with 1,522 in August 1997. The rate of natural increase went down from 3.3 to 2.9 per 1,000 population during this period.

**Marriages** decreased for all three time periods shown below while **dissolutions of marriage** increased for all three periods. The marriage to divorce ratio for the 12 months ending with August decreased from 1.77 to 1.62.

**Infant deaths** decreased in August as 40 Missouri babies died compared with 43 one year earlier. The infant death rate for January-August increased from 8.0 to 8.4 per 1,000 live births.

### PROVISIONAL RESIDENT VITAL STATISTICS FOR THE STATE OF MISSOURI

Item	August				Jan.-Aug. cumulative				12 months ending with August				
	Number		Rate*		Number		Rate*		Number		Rate*		
	1997	1998	1997	1998	1997	1998	1997	1998	1997	1998	1996	1997	1998
<b>Live Births .....</b>	5,750	5,908	12.5	12.4	49,143	48,947	13.7	13.5	72,842	74,385	13.6	13.5	13.7
<b>Deaths .....</b>	4,228	4,530	9.2	9.5	37,245	35,971	10.4	9.9	54,761	53,564	10.1	10.2	9.9
<b>Natural increase ....</b>	1,522	1,378	3.3	2.9	11,898	12,976	3.3	3.6	18,081	20,821	3.5	3.4	3.8
<b>Marriages .....</b>	4,773	4,754	10.4	10.0	29,836	28,278	8.3	7.8	44,717	42,254	8.4	8.3	7.8
<b>Dissolutions .....</b>	2,005	2,235	4.4	4.7	16,991	17,471	4.7	4.8	25,326	26,100	4.8	4.7	4.8
<b>Infant deaths .....</b>	43	40	7.5	6.8	395	411	8.0	8.4	600	584	7.4	8.2	7.9
<b>Population base .....</b> (in thousands)	...	...	5,402	5,440	...	...	5,402	5,440	...	...	5,351	5,389	5,427

\* Rates for live births, deaths, natural increase, marriages and dissolutions are computed on the number per 1000 estimated population. The infant death rate is based on the number of infant deaths per 1000 live births. Rates are adjusted to account for varying lengths of monthly reporting periods.

AN EQUAL OPPORTUNITY/AFFIRMATIVE ACTION EMPLOYER - Services provided on a nondiscriminatory basis.

Alternate forms of this publication for persons with disabilities may be obtained by contacting the Missouri Department of Health, Center for Health Information Management & Epidemiology/Bureau of Health Data Analysis, P.O. Box 570, Jefferson City, MO 65102; phone (573) 751-6278. Hearing-impaired citizens telephone 1-800-735-2966.